

Paper March 1828

An
Inaugural Essay
on
Pertussis,
for

The degree of M. D.

In

The University of Pennsylvania:

By

William M. Fenimore

Of

New Jersey.

June 5th 1828.

Market Street

295

227. *Spiz. mont.* 1822

Chrysomitris 1822

Spiz. mont.

Spiz. mont.

Spiz. mont.

Spiz. mont.

Spiz. mont.

Spiz. mont.

Spiz. mont.

Spiz. mont.

Spiz. mont.

228

Peritussis, or whooping cough, which I have selected for the subject of this dissertation, is a disease, which is, almost exclusively, confined to the ages of infancy, and childhood; and, which, having once run its course, commonly, exonerates the system, from the liability to a second attack: Yet, numerous instances are recorded, as well, of its having attacked persons at an advanced period of life, as, of its having occurred repeatedly in the same individuals.

An attack of this disease is, usually, ushered in with the ordinary symptoms of catarrh. — There is a sense of lachrymation, & weakness, with pain in the head; some degree of hoarseness, dry cough; frequent sneezing, with a discharge of thin mucus from the nostrils; loss of appetite, and, commonly, more or less febrile excitement: — These symptoms, frequently, continue for many days, before the disease appears in its usual form. And, it has, even, been said by Cullen, that, "he has known instances of a disease, which, though evidently arising from the chris cough contagion, never put on any other form, than that of common catarrh." But this does not often happen, and, more generally,

in the co
clandestine
something.
I forgot
following
continued,
apparently,
by a violen
ance, call
the popula
opinions
narrow, by
you contain
from the la
by omitting
of temporary
common, some
was, unlike
brought to the

in the course of two or three weeks, the disease takes on its characteristic symptoms, particularly, the paroxysm of convulsive coughing.

A paroxysm consists in a number of short convulsive expirations, following each other in rapid succession; and, which are sometimes continued, until the patient becomes nearly exhausted, and is, apparently, on the point of suffocation, when the lungs are filled, by a violent inspiration, which is attended by that peculiar noise, called, a whoop; (from whence, the disease has received the popular appellation of whooping cough.) — the convulsive expirations are now renewed, and, again, terminated in the same manner, by the powerful inspiration; and, in this way, the paroxysm continues, until, there is a discharge of phlegm or mucus, from the lungs, or, until the contents of the stomach are ejected by vomiting; after which, with the exception, of some degree of temporary exhaustion, and uneasiness of the chest, the child, in common, remains cheerful and lively, and, apparently, free from disease, until the coming on of the next paroxysm, when, an, again, presented the same phenomena as above described.

[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]

After the
strong de
low vacu
long a p
the long, is
knowledge
the lips are
the socks
there are a
to some of
suffer, for
a person
himself on
until it is
ful of the
then the co
the felt of
sometimes, or
no time, in

After the termination of the paroxysm, the patient has, commonly, a strong desire for food, and, more especially, if the stomach has been evacuated by vomiting.

During a paroxysm, the blood, from its interrupted circulation thro' the lungs, is, sometimes, forced to the head with such violence, that, hemorrhage from the nose, does not unfrequently take place; the lips and eyelids swell; the eyes appear ready to start from their sockets; the countenance becomes of a livid hue; and these are all the symptoms of immediate strangulation:

A sense of dizziness, also, arises from the same cause, and the sufferer, from a consciousness of this, at the commencement of a paroxysm, clings to the nearest object for support, or throws himself on his hands and knees, and in this position remains, untill its violence has abated. — At other times, the ef-

fects of the fit are much milder, and during its continuation, the child experiences little inconvenience. —

The fit of coughing occurs with various degrees of frequency; sometimes, in mild cases, not happening, oftener, than five or six times, in the course of twenty-four hours; while, in more vi-

[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]

but all
may be
as you
perhaps
to itself
for ever
to cover
will dep
as relate
an other
brasted
He to
of opinion
while it
when it
from it
knowledge
where
the ad

lent attacks, they may occur every half hour, especially during the night, when, they are always more frequent, as well as, more severe.

Portupis, being once confirmed, frequently continues, if left to itself, for three or four months, and sometimes, tho' rarely for over ten or twelve months; but, when judiciously treated, its course may be considerably abbreviated. Much, however, will depend on the season of the year in which it occurs, both as relates to its duration, and the severity of its symptoms, an attack being always more violent, and its course more protracted during winter.

As to the origin of portupis, there is not a little difference of opinion: many firmly believing in its contagious nature, while others, entirely, deny its propagation in that manner, and ascribe its production altogether to atmospherical influences; and, from its frequent prevalence as an epidemic, together, with the knowledge of its having, occasionally, broken out in situations where there had been no intercourse with an infected source, the advocates of its non-contagious nature appear to be entitled

[Faint, illegible handwriting on the left page of an open notebook. The text appears to be a continuous paragraph across approximately 15 lines.]

to the great
it is his
either on
beginning
can be
mean. 7
on the re
could it be
participate
has declar
then has
effect of
a pretty g
natural, a
of super
nature de
sympath
obstruction
nature m

to the greater share of credit.

Now, is there less diversity of sentiment among those who have written on this disease, with respect to their pathological view, disagreeing, as well, in the organ in which they consider the disease to be primarily located, as, in the nature of its proximate cause. The opinion has been advanced by some, that the brain was the original seat of the disorder: while others have located it in the alimentary canal: the lungs being said to participate in the disease, merely, from sympathy: Some have declared it to be, entirely, of a spasmodic nature, while others have described its phenomena, as being, exclusively, the effects of inflammation. But, the notion, which at present is pretty generally entertained, and which appears the most rational, is, that pertussis is a primary affection of the organs of respiration: any other organs or tissues that may become involved during the course of the disease, be so, either from sympathetical influence, or from the effects of mechanical obstructions during the paroxysms. — The remote cause, whatever may be its nature, is applied to the mucous membrane,

[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]

which has
of degree
forming a
dion of
les, and
unduly
described
known
matters in
consequence
has char
ness, a
will be for
the deep
the broken
tapes, in
frequently
dysplasia,
from the

which lines the trachea and bronchioles, exciting a greater or less degree of irritation, strongly tending to, and frequently, running into inflammation; as a consequence of this condition of the secreting membrane, the fluids become irritated, and increased in quantity; and, for their expulsion, when unduly accumulated, the paroxysms of coughing, as above described, are called into action. Fits of coughing may likewise be excited by various causes, as the irritation of acid matters inhaled with the breath, violent exercise, laughter &c. Concerning the diagnosis, not much need be said. The peculiar character of the cough, considered with the more gradual manner, in which an attack of pertussis makes its approach, will be found sufficient, to distinguish it from other affections of the respiring organs; as Spasmodic Asthma, Bronchitis, Cynanche trachealis &c. There is, however, a disease, to which pertussis, in its first stage, bears so strong a resemblance, that, frequently, if we confine our attention entirely to the existing symptoms, it will be found no easy matter, to distinguish the one from the other. — I allude to an attack of common cold

for cold.
 plates wait
 for an other
 part, in case
 will be an
 volume of
 to find, &
 day, of a
 subject. It
 only, be a
 to provide
 our judgment
 mistake, as
 a possibly
 to propagate
 of this disc
 its consider
 the evidence
 situation.

from cold. There is, however, in the latter, commonly, more febrile excitement, and the patient is less disposed to coma than in whooping cough. But, the most certain diagnosis will consist, in our knowledge of the prevailing disease: — for, if a child be attacked with catarrhal symptoms, during the prevalence of whooping-cough as an epidemic, it will, generally, be found, that these, are, merely, the forerunners of other symptoms, of a character so peculiar, as to leave no doubts on the subject. And, under these circumstances, we will, most commonly, be correct, in deciding, the disease to be an attack of the prevailing epidemic; and, should we ever be wrong in our judgement, little importance need be attached to the mistake, as, at this period, the treatment to be pursued, is precisely similar in either case.

To prognosticate with any degree of certainty, how an attack of this disease will terminate, it will be necessary to take into consideration, the period of life at which it takes place, the violence of the attack, and the state of the patient's constitution.

He says,
 greater a
 it grows
 say, than
 the more
 told to b
 say every
 by Dr. G
 say of the
 struggle
 or those
 founders
 that a be
 the instab
 we cons
 large by
 larger of
 greater a

As to the time of life, at which, an attack takes place, Dr. Caldwell says, "the younger that children are, they are in the greatest danger from this disease, and of those to whom it proves fatal, there are many more under two years of age, than above it: And the older children are, they are the more secure against an unhappy event; and this I hold to be a very general rule, though, I own, there are many exceptions to it." But, on the other hand, it is affirmed by Dr. Gregory, that, "the danger is not proportioned to the age of the child, for, one, of two or three months old, will struggle through the disease, as well as, another, of two or three years." This latter opinion is, perhaps, not founded on accurate observation, as, it is generally found that a very tender age is unfavourable, principally, from the inability of infants to obtain relief by expectoration and consequently, unless the mucus is expelled from the lungs by the action of vomiting, they are, sometimes, in danger of being suffocated by it; and, hence arises the greater utility of emetics, in their management.

The principal
as disease
of broncho
ing, all the
trachea is
given her
from the
of an allu
valve, ins
the state o
taken into
be improv
her debilit
infantum,
the intes
an improv
children
to pulmon
or an very

The principal symptoms that denote a severe and dangerous disease, are much febrile excitement, great difficulty of breathing, and violent and frequent paroxysms of coughing, attended with scanty expectoration: copious expectoration in very young infants, is likewise unfavourable: copious hemorrhage from the nose, are generally, injurious, from the debility they occasion: The super-
 vention of an attack of pleurisy, or other inflammatory disease, is, also, inauspicious, and frequently, fatal.

The state of the patients constitution is, likewise, to be taken into consideration, in forming a prognosis. It will be unfavourable, if the attack occur in those, who have been debilitated by some previous disease, as cholera infantum, diarrhoea &c.

The irritation, produced by dentition, will, likewise, have an unfavourable effect on the disease.

Children who inherit from their parents, predilections to pulmonary affections, as *phthisis pulmonalis*, asthma, &c. are very obnoxious to a fatal terminating end, and, are,

induced,
topis o
no inf
nysmo
a free
during
to sleep
ficially
-induced
in dec
particip
Life and
by suff
salvatio
system;
being in
to the
-phony, c
Loudon

indeed, seldom known to recover. But, when pertussis occurs to children of sound constitutions, — when no inflammatory affection supervenes, — when the paroxysms are neither frequent, nor violent, and terminate in free expectoration, or vomiting, — and, when the patient, during the intermissions, has a desire for food, a disposition to sleep, and, is neither, troubled with fever, nor difficultly of breathing, the case may, commonly, be considered, as without danger, and, we will be justifiable in delivering a favourable prognosis.

Pertussis when fatal, may terminate in various ways.

Life may be, instantaneously, destroyed, during the paroxysm, by suffocation; or the patient may be carried off by convulsion, brought on by the violent shock received by the system; or, the free transpiration of blood through the lungs, being interrupted, the consequent increased determination to the vessels of the brain, may produce an attack of apoplexy, immediately destructive of life.

Foundation may be laid for other diseases, which may

form, a
 narrow
 flange,
 in the
 flange
 water,
 galapier
 remains,
 but, for
 lack of
 sing out
 poor.
 Treatments
 of disease
 of equal
 modes of
 care, in the
 employment
 in cases

prove, ultimately, fatal, though in a much more gradual manner: thus, from the extension of inflammation to the pleura, serous effusions may be produced, constituting hydrothorax; or, the tunica arachnoidea may take on inflammatory action, coagulating in hydrocephalus internus:

Weitz, likewise, is said, sometimes, to follow an attack of pertussis. — Pleurisy palmonalis is not an unrequent occurrence, especially, in those, who are predisposed to it. — But, perhaps, the most common manner, in which, an attack of pertussis prove fatal to life, is, by gradually wearing out the strength of the patient, by marasmus, or hectic fever. — *quædam agunt the effects of it, in order to give*

Treatment. It is very probable, that in the whole catalogue of diseases to which the human race is subject, there is not one of equal severity, that, in general, is subjected to more empirical modes of treatment, than whooping cough; Nor, perhaps, is there one, in which, more benefit may, occasionally, be derived, from the employment of appropriate remedies.

In cases which are very mild, and, unaccompanied with febrile

customer
exist,
apostrophe
by the
between
liquidity
simply
in order
or, the
position
these
can't
went to
attending
But, I
a more
its poor,
of the
be paid

evilemias, or dyspnoea, and, in which the disease appears to consist, chiefly, in the paroxysm of coughing, little medical assistance, indeed, is requisite. Emollients should be avoided, by the occasional use of castor oil, or rhubarb. — Small doses of Nalimomial tincture, or, what is better perhaps, the Comp. Symp. of Squills, may, also, be employed, to facilitate expectoration, occasionally (especially in very young infants) increasing the dose, in order to produce vomiting. Antispasmodics, as Asafetida, or, the tincture of Musks, are useful, to allay the cough; and, sometimes, will speedily arrest the progress of the disease. These means, together with attention to diet, and taking care to guard against the effects of cold, in order to prevent the lightening up of inflammation, will, commonly, be all-sufficient.

But, should the attack, from the commencement, be of a more violent character, or, should it become so during its progress, it will be highly necessary for the welfare of the patient, that a greater share of attention should be paid to the disease, and its violence combatted by

were offered
 to local
 education
 and for
 operating
 to all
 a full
 have been
 -telling is
 their
 to the
 should
 should be
 best
 from the
 case, which
 is of a
 dispersed
 parties

more effectual remedies.

In treating the disease, under these circumstances, the first indication evidently is to moderate the violence of the disease, and palliate the urgent symptoms: And, for this purpose, the evacuating remedies are undoubtedly to be employed.

In all such cases, attended with much febrile excitement, with a full hard pulse, and great difficulty of breathing, or, if there be symptoms of pneumonia inflammation, general blood-letting is indispensable, particularly, in those of full plethoric habits: The quantity to be drawn is to be proportioned to the age of the patient, and the severity of the symptoms, and, should relief not be obtained by the first bleeding, the operation should be repeated.

Much benefit will also be derived, by detracting blood locally from the chest, by means of cups or leeches; and, in many cases, when the symptoms are less violent, or, when the patient is of a very tender age, general bleeding may be entirely dispensed with, and, topical depletion alone, considered in.

Emetics are also to be employed, and, the advantage to be

the device
 is relative
 justice a
 If the
 have been
 single on
 motives;
 applicable
 treated a
 upon
 auxiliary
 able to c
 appear in
 not so we
 If this cla
 employed,
 efficiency,
 administration
 It may

be derived from this an. may be inferred, from what is said in relation to them, by Dr Chapman, in his treatise on *Prætica Medica and Therapeutica*.

Of the various remedies, "say Si", that at different periods have been suggested for the cure of ptochus, I know of no single one from which I have derived better effects, than from emetics. To the earlier stage of the complaint they are chiefly applicable, and when the attack is violent, they must be repeated daily, or even twice a day, for a week or two in succession. By steadily persisting in this course, and with the auxiliary means proper to the case, we shall generally be able to conduct the case to a comparatively speedy and successful issue, provided it be that of a child, as the remedy is not so well adapted to persons in more advanced life.))

Of this class of remedies, the tartarized antimony is generally employed, being preferred, as well, on account of its superior efficacy, as, from the greater facility with which it may be administered to children, than most other articles of the class. It may be given, either, in the form of Antimonial wine,

on that
 to go to
 day to co
 add
 line of 3
 with spray
 after the
 like effec
 the comp
 the spray
 day, the
 increase
 be subst
 Pangeles
 be condit
 they are a
 bad whe
 the board
 administr

or in that of watery solution. Of the wine, a child within the year may take ten, fifteen, or twenty drops, or more, according to circumstance, at short intervals, until vomiting is excited. The watery solution may be prescribed in the proportion of 3 grains of the Antimony, to 6 ounces of water, with a little Syrup. Of this, a large teaspoonful may be given every fifteen minutes, to a child of one or two years old, until it take effect.

The comp. Syrup of Squills, or, as it is more commonly called, the "Nise Syrup," is also an invaluable article. To produce vomiting, the dose should be large and frequently repeated.

Specacchara, Sulphate of Zinc, and other articles, may likewise be substituted for the tartarised antimony.

Purgative medicines are also demanded, on account of the costive condition of the bowels, which commonly attends the disease. They are also serviceable, by relieving those determinations to the head which so frequently exist.

The bowels should be evacuated, every two, or three days, by the administration of a mercurial purge. A child of a year old

may take
polish
spicily
little can
may also
a solution
after 10
of bread
the apple
dried
Expecta
great be
squalid
The appa
by the m
inegar, or
After the
the infla
ious on

may take 2 or 3 grains of calomel, in combination with a small portion of jalap or rhubarb; and, if the bowels be not sufficiently opened by this, the operation may be assisted by a little castor oil. — Rhubarb alone, or an infusion of senna, may also be employed with advantage, to keep the bowels in a soluble condition. *Use of a gunn.*

After bleeding, and other evacuating remedies, if the difficulty of breathing, and sense of oppression, still remain unsatisfied, the application of a blister to the chest will be attended with decided advantage. *the marked sensibility of the*

Expectorant medicines of various kinds are sometimes of great benefit: as speccacunda, antimonial wine, syrup of squills &c. in small doses, frequently repeated. —

The expectoration of the viscid phlegm will also be facilitated, by the inhalation of various vapours; as, of warm water & vinegar, or ether &c. The turpentine is also useful. —

After the violence of the disease has been moderated, and, the inflammatory symptoms have been subdued, by the various remedies employed for that purpose, Articles of an

nearly a
 new g
 Be fixed
 -ed in
 of appar
 Since a
 Nearly a
 son, pro
 direction
 awful
 fested pa
 of them p
 awful:
 Digitalis
 to be of m
 to in the f
 a child of
 female of
 every day

entirely different character are to be resorted to, with a view of arresting the progress of the disease.

The fixed alkalies are much employed. They are prescribed in doses of 2 or 3 grains, with a few drops of wine of Spicaeantha and one drop of Laudanum, two or three times a day to a child of a year old.

Nearly all the articles of the class of Narcotics, are likewise prescribed, at different times, and after the reduction of the more violent symptoms, they may be useful by allaying the morbid sensibility of the affected parts, and thereby putting a stop to the disease. Of these, perhaps, Opium will be found, in common, the most useful. The best mode of prescribing it is in watery solution. Digitalis is said by Dr Pearson, in his treatise on Chincough, to be of most decided efficacy; his mode of prescribing it is in the form of saturated tincture, of which, he says a child of a year old may safely take 6 drops, at intervals of six hours; the dose to be increased, one drop every day, untill a diuretic effect is produced, or untill

is often
Killock
is far
not un-
Adverse
phymen
Tension
by the
the train
they fear
the ton
particu-
not em-
inflam-
injurious,
Not a
interval,
-case. It
a little

its effect on the pulse, demands its suspension.

Henlock, Prussic acid, Belladonna, Stramonium &c are all frequently administered with the same view, and not without occasional benefit.

Advantage is likewise to be obtained from the employment of Antispasmodics, as *Opipetida*, *Musk* &c. Sometimes, the disease appears to be prolonged merely by the force of habit, and in this case, to break up the train of morbid affections, and restore the healthy functions of the system, the peruvian bark and other Tonics appear to be peculiarly appropriate, but particular care should be taken that these articles be not employed, while there is the slightest evidence of inflammation, or their administration will be decidedly injurious.

Not a small proportion of praise is conferred on the internal use of cathartics, at this period of the disease. It is commonly prescribed in tincture, with a little Laudanum; commencing with small doses

as produced
 after occur-
 ing exactly
 Simola's
 of certain
 given, and
 have a go
 to change
 after all
 that atten-
 the whole c
 treatment is
 of practice
 performed
 article is
 risk is spec-
 to be emp-
 conclusion
 to be seen

and gradually increasing them untill strangury is induced; on the occurrence of which, the progress of the disease is frequently arrested.

Stimulating articles, as spirits of turpentine, tincture of cantharide, juice of garlic, &c rubbed along the spine, and applied to the soles of the feet, sometimes have a good effect.

A change of residence will sometimes perform a cure, after all other means have failed.

Strict attention to diet is highly necessary throughout the whole course of the disease, and the success of the treatment will depend in a great measure, on the degree of exactness with which the directions in this respect are performed: it should consist of light and unirritating articles in the first stage, but, after the force of the attack is spent, those of a more nutritive character are to be employed.

Convulsions sometimes occur during the paroxysms; they are to be relieved by the use of the warm bath, the

applicat
Hydroth
regula
rest, of
-part is a

application of leeches to the temple, purging &c. —

Hydrothorax, Hydrocephalus, marasmus, and the various
sequela of the disease, require peculiar form of treat-
ment, of which nothing is to be said here, as the sub-
-ject is of a nature, unconnected with this discussion.

